


IgA Antibody Investigation Request

Instructions for filling this form. The form can be filled in using your PC:

1. Complete appropriate sections on Page 1.
2. Provide required additional information as indicated on Page 2.
3. Collect the appropriate sample tubes as specified on Page 2.
4. Send samples with a completed copy of this request form.

Testing Laboratory	QLD (QTIS) Platelet and Neutrophil Reference Laboratory		
Please send samples to	Platelet and Neutrophil Reference Laboratory Australian Red Cross Lifeblood 44 Musk Avenue, (deliveries via Blamey Street), Kelvin Grove, Brisbane, QLD 4059		
Phone	(07) 3838 9173 or 38383 9487	Email	DL- AUSPlateletandNeutrophil@redcrossblood.org.au

Patient details				
Last Name		First Name		
Gender		MRN/UR	DOB	
Referring Clinician details				
Name			Date requested	
Signature			Phone	
Address <i>Tick if a hard copy report is required</i>			Email	
Referring Laboratory name				
Phone		Email		
Name of person completing the form (if different from above)				
Name				
Phone		Email		
Sample collection				
Collector's name		Date & time of collection		
Patient's signature		Date		
Specimen type	Serum (clot)			
Please attach label/barcode 				

Patient details					
Last Name			First Name		
Gender		MRN/UR		DOB	
Clinical details					

IgA Antibody – Indication for testing and additional clinical information as required

IgA Antibody Testing		
IgA Level	g/L	
Date tested		
Has this patient received blood products in the past?	Yes	No
Any history of anaphylactic reactions to blood products?	Yes	No
If Yes, date of transfusion/reaction		
Details of transfusion/reactions		

IgA Antibody Sample Collection Guidelines

Investigation request and samples	Special instructions and indicative turnaround time	Storage and transport instructions				
IgA deficient patient <table border="0"> <tr> <td>Adult</td> <td>12 mL serum (clot)</td> </tr> <tr> <td>Child</td> <td>2 - 4 mL serum (clot)</td> </tr> </table>	Adult	12 mL serum (clot)	Child	2 - 4 mL serum (clot)	No special instructions <i>Note: Laboratory turnaround time is 3 working days.</i>	Store and transport at either room temperature or 4°C.
Adult	12 mL serum (clot)					
Child	2 - 4 mL serum (clot)					
IgA antibodies for anaphylactic reactions <table border="0"> <tr> <td>Adult</td> <td>12 mL serum (clot)</td> </tr> <tr> <td>Child</td> <td>2 - 4 mL serum (clot)</td> </tr> </table>	Adult	12 mL serum (clot)	Child	2 - 4 mL serum (clot)	Please send both pre and post transfusion samples <i>Note: Laboratory turnaround time is 3 working days.</i>	
Adult	12 mL serum (clot)					
Child	2 - 4 mL serum (clot)					